NINDS CDE Notice of Copyright Epworth Sleepiness Scale (ESS) – Adult Version

| Availability: | Please visit this website for the instrument: |
|---------------------------------|--|
| | Epworth Sleepiness Scale. |
| Classification: | Supplemental – Highly Recommended: Spinal Cord Injury (SCI)-Pediatric |
| | Supplemental: Sports-Related Concussion (SRC) Subacute (after 72 hours to 3 months), and Persistent/Chronic (3 months and greater post concussion) |
| | Exploratory: Sports-Related Concussion (SRC) Acute (time of injury until 72 hours) |
| Short Description of Instrument | The Epworth Sleepiness Scale (ESS) developed in 1991 is the most widely used subjective scale of daytime somnolence. The current version used is the 1997 version. |
| | As with all subjective scales, participants may not recognize sleepiness, particularly when chronic. |
| | Translations are available in over 20 languages and the Scale can be completed in less than 5 minutes., one needs a license to use it. |
| Scoring: | Rated from 0–24, with high scores indicating worse sleepiness. |
| | General interpretation is as follows |
| | 0–5 Lower Normal Daytime Sleepiness |
| | 6–10 Higher Normal Daytime Sleepiness |
| | 11–12 Mild Excessive Daytime Sleepiness |
| | 13–15 Moderate Excessive Daytime Sleepiness |
| | 16–24 Severe Excessive Daytime Sleepiness |
| References: | Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. Sleep. 1991;14(6):540–545. |
| | Johns MW. A new perspective on sleepiness. Sleep Biol Rhythm, 2010; 8: 170-179. |
| | Melendres MC, Lutz JM, Rubin ED, Marcus CL. Daytime sleepiness and hyperactivity in children with suspected sleep-disordered breathing. Pediatrics. 2004;114(3):768–775. |
| | Meltzer LJ, Mindell JA. Sleep and sleep disorders in children and adolescents. Psychiatr Clin North Am. 2006;29(4):1059–1076. |
| | Mollayeva T, Shapiro CM, Cassidy JD, Mollayeva S, Colantonio A. Assessment of Concussion/Mild Traumatic Brain Injury-Related Fatigue, Alertness, and Daytime Sleepiness: A Diagnostic Modelling Study. Neuropsychiatry (London) 6.6 (2016): 525–543. |